

SAMPLE OUTPUT FROM 'polyinterp'

```
> res1 := polyinterp([ [0,1], [1,6], [2,4], [3,0] ],'x');
```

$$res1 := \frac{5}{6}x^3 - 6x^2 + \frac{61}{6}x + 1$$

```
> [subs(x=0,res1), subs(x=1,res1),subs(x=2,res1), subs(x=3,res1)];
```

$$[1, 6, 4, 0]$$

```
> res2:= polyinterp([ [0,1], [1,6], [2,4], [3,0] ],f(x));
```

```
Error, (in polyinterp) second argument must be a name
```

```
> res3 := polyinterp( [ [ -h , f[-1] ], [ 0 , f[0] ], [ h , f[1] ] ] , 'x' );
```

$$res3 := \frac{1}{2} \frac{f_{-1} x^2}{h^2} - \frac{1}{2} \frac{f_{-1} x}{h} - \frac{f_0 x^2}{h^2} + f_0 + \frac{1}{2} \frac{f_1 x^2}{h^2} + \frac{1}{2} \frac{f_1 x}{h}$$

```
> res3c := collect(res3,{f[-1],f[0],f[1]});
```

$$res3c := \left(\frac{1}{2} \frac{x^2}{h^2} - \frac{1}{2} \frac{x}{h} \right) f_{-1} + \left(- \frac{x^2}{h^2} + 1 \right) f_0 + \left(\frac{1}{2} \frac{x^2}{h^2} + \frac{1}{2} \frac{x}{h} \right) f_1$$