PHYS 410/555 Computational Physics

Solution of Non Linear Equations (a.k.a. Root Finding)

(Reference *Numerical Recipes*, 9.0, 9.1, 9.4)

- We will consider two cases

1. $f(x) = 0$ "1-dimensional"
2. $\mathbf{f}(\mathbf{x}) = \mathbf{0}$ "d-dimensional"

$$\mathbf{x} \equiv [x_1, x_2, \ldots, x_d]$$

$$\mathbf{f} \equiv [f_1(x_1, x_2, \ldots, x_d), \ldots, f_d(x_1, x_2, \ldots, x_d)]$$

1. Solving Nonlinear Equations in One Variable

- We have briefly discussed bisection (binary search), will consider one other technique: *Newton's method* (Newton-Raphson method).

Preliminaries

- We want to find one or more roots of

$$f(x) = 0 \tag{1}$$

We first note that *any* nonlinear equation in one unknown can be cast in this canonical form.

- *Definition:* Given a canonical equation, $f(x) = 0$, the *residual* of the equation for a given x-value is simply the function f evaluated at that value.

- *Iterative technique:* Assume $f(x) = 0$ has a root at $x = x^\star$; then consider sequence of estimates (iterates) of $x^\star$, $x^{(n)}$

$$x^{(0)} \rightarrow x^{(1)} \rightarrow x^{(2)} \rightarrow \cdots \rightarrow x^{(n)} \rightarrow x^{(n+1)} \rightarrow \cdots \rightarrow x^\star$$

- Associated with the $x^{(n)}$ are the corresponding residuals

$$\begin{array}{ccccccccccc} r^{(0)} & \rightarrow & r^{(1)} & \rightarrow & \cdots & \rightarrow & r^{(n)} & \rightarrow & r^{(n+1)} & \rightarrow \cdots \rightarrow & 0 \\ \| & & \| & & & & \| & & \| & & \| \\ f(x^{(0)}) & & f(x^{(1)}) & & & & f(x^{(n)}) & & f(x^{(n+1)}) & & f(x^\star) \end{array}$$

Locating a root $\equiv$ Driving the residual to 0

- *Convergence:* When we use an iterative technique, we have to decide *when* to stop the iteration. For root finding case, it is natural to stop when

$$|\delta x^{(n)}| \equiv |x^{(n+1)} - x^{(n)}| \leq \epsilon \qquad (2)$$

where ϵ is a prescribed convergence criterion.

- A better idea is to use a "relativized" δx

$$\frac{|\delta x^{(n)}|}{|x^{(n+1)}|} \leq \epsilon \tag{3}$$

but we should "switch over" to "absolute" form (2) if $|x^{(n+1)}|$ becomes "too small" (examples in on-line code).

- *Motivation:* Small numbers often arise from "unstable processes" (numerically sensitive), e.g. $f(x+h) - f(x)$ as $h \to 0$, or from "zero crossings" in periodic solutions etc.—in such cases may not be possible and/or sensible to achieve stringent relative convergence criterion

Newton's method

- Requires "good" initial guess, $x^{(0)}$; "good" depends on specific nonlinear equation being solved
- Refer to Numerical Recipes for more discussion; we will assume that we have a good $x^{(0)}$, and will discuss one general technique for determining good initial estimate later.
- First consider a rather circuitous way of solving the "trivial" equation

$$ax = b \quad \longrightarrow \quad f(x) = ax - b = 0 \tag{4}$$

Clearly, $f(x) = 0$ has the root

$$x^\star = \frac{b}{a} \tag{5}$$

- Consider, instead, starting with some initial guess, $x^{(0)}$, with residual

$$r^{(0)} \equiv f(x^{(0)}) \equiv ax^{(0)} - b \tag{6}$$

Then we can compute an improved estimate, $x^{(1)}$, which is actually the solution, $x^\star$, via

$$x^{(1)} = x^{(0)} - \delta x^{(0)} = x^{(0)} - \frac{r^{(0)}}{f'(x^{(0)})} = x^{(0)} - \frac{f(x^{(0)})}{f'(x^{(0)})} \tag{7}$$

"Proof":

$$x^{(1)} = x^{(0)} - \frac{r^{(0)}}{a} = x^{(0)} - \frac{ax^{(0)} - b}{a} = \frac{b}{a} \tag{8}$$

- Graphically, we have



- *Summary*

$$x^{(1)} = x^{(0)} - \delta x^{(0)} \tag{9}$$

where $\delta x^{(0)}$ satisfies

$$f'(x^{(0)})\delta x^{(0)} = f(x^{(0)}) \tag{10}$$

or

$$f'(x^{(0)})\delta x^{(0)} = r^{(0)} \tag{11}$$

- Equations (9-10) immediately generalize to non-linear $f(x)$ and, in fact, are precisely Newton's method.

• For a general nonlinear functrion, $f(x)$, we have, graphically



- *Newton's method for* $f(x) = 0$*:* Starting from some initial guess $x^{(0)}$, generate iterates $x^{(n+1)}$ via

$$x^{(n+1)} = x^{(n)} - \delta x^{(n)} \tag{12}$$

$$f'(x^{(n)})\delta x^{(n)} = r^{(n)} \equiv f(x^{(n)}) \tag{13}$$

or more compactly

$$x^{(n+1)} = x^{(n)} - \frac{f(x^{(n)})}{f'(x^{(n)})} \tag{14}$$

- *Convergence: When* Newton's method converges, it does so *rapidly*; expect number of significant digits (accurate digits) in $x^{(n)}$ to roughly *double* at each iteration (quadratic convergence)

- *Example: "Square Roots"*

$$f(x) = x^2 - a = 0 \quad \longrightarrow \quad x^\star = \sqrt{a} \tag{15}$$

Application of (14) yields

$$\begin{aligned} x^{(n+1)} &= x^{(n)} - \frac{{x^{(n)}}^2 - a}{2x^{(n)}} \\ &= \frac{2{x^{(n)}}^2 - \left({x^{(n)}}^2 - a\right)}{2x^{(n)}} \\ &= \frac{{x^{(n)}}^2 + a}{2x^{(n)}} \end{aligned}$$

which we can write as

$$x^{(n+1)} = \frac{1}{2}\left(x^{(n)} + \frac{a}{x^{(n)}}\right) \tag{16}$$

- Try it manually, compute $\sqrt{2} = 1.414\ \ 2135\ \ 6237$ using 12-digit arithmetic (hand calculator)

Iterate	Sig. Figs
$x^{(0)} = 1.\mathbf{5}$	1
$x^{(1)} = \frac{1}{2}\left(1.5 + 2.0/1.5\right) = 1.41\mathbf{6}\ \ 6666\ \ 6667$	3
$x^{(2)} = \frac{1}{2}\left(1.416\cdots + 2.0/1.416\cdots\right) = 1.414\ \ 21\mathbf{5}6\ \ 8628$	6
$x^{(3)} = \frac{1}{2}\left(1.4142\cdots + 2.0/1.4142\cdots\right) = 1.414\ \ 2135\ \ 623\mathbf{8}$	11

Alternate Derivation of Newton's Method (Taylor series)

- Again, let $x^\star$ for a root of $f(x) = 0$, then

$$0 = f(x^\star) = f(x^{(n)}) + (x^\star - x^{(n)})f'(x^{(n)}) + O((x^\star - x^{(n)})^2) \quad (17)$$

Neglecting the higher order terms, we have

$$0 \approx f(x^{(n)}) + (x^\star - x^{(n)})f'(x^{(n)}) \quad (18)$$

Now, treating the last expression as an equation, and replacing $x^{(n)}$ with the new iterate, $x^{(n+1)}$, we obtain

$$0 = f(x^{(n)}) + (x^{(n+1)} - x^{(n)})f'(x^{(n)}) \quad (19)$$

or

$$x^{(n+1)} = x^{(n)} - \frac{f(x^{(n)})}{f'(x^{(n)})} \quad (20)$$

as previously.

2. Newton's Method for Systems of Equations

- We now want to solve

$$\mathbf{f}(\mathbf{x}) = \mathbf{0} \tag{21}$$

where

$$\mathbf{x} = (x_1, x_2, \ldots, x_d) \tag{22}$$

$$\mathbf{f} = (f_1(\mathbf{x}), f_2(\mathbf{x}), \ldots, f_d(\mathbf{x})) \tag{23}$$

- *Example ($d = 2$):*

$$\sin(xy) = \frac{1}{2} \tag{24}$$

$$y^2 = 6x + 2 \tag{25}$$

In terms of our canonical notation, we have

$$\mathbf{x} \equiv (x, y) \tag{26}$$

$$\mathbf{f} \equiv (f_1(\mathbf{x}), f_2(\mathbf{x})) \tag{27}$$

$$f_1(\mathbf{x}) = f_1(x, y) = \sin(xy) - \frac{1}{2} \tag{28}$$

$$f_2(\mathbf{x}) = f_2(x, y) = y^2 - 6x - 2 \tag{29}$$

- The method is again iterative, we start with some initial guess, $\mathbf{x}^{(0)}$, then generate iterates

$$\mathbf{x}^{(0)} \to \mathbf{x}^{(1)} \to \mathbf{x}^{(2)} \to \cdots \to \mathbf{x}^{(n)} \to \mathbf{x}^{(n+1)} \to \cdots \to \mathbf{x}^{\star}$$

  where $\mathbf{x}^{\star}$ is *a* solution of (21)

- *Note:* The task of determining a good initial estimate $\mathbf{x}^{(0)}$ in the d-dimensional case is even more complicated than it is for the case of a single equation—again we will *assume* that $\mathbf{x}^{(0)}$ *is* a good initial guess, and that $\mathbf{f}(\mathbf{x})$ is sufficiently well-behaved that Newton's method will provide a solution (i.e. *will* converge).

- As we did with the scalar (1-d) case, with any estimate, $\mathbf{x}^{(n)}$, we associate the *residual vector*, $\mathbf{r}^{(n)}$, defined by

$$\mathbf{r}^{(n)} \equiv \mathbf{f}(\mathbf{x}^{(n)}) \tag{30}$$

- The analogue of $f'(x)$ in this case is the *Jacobian matrix*, $\mathbf{J}$, of first derivatives. Specifically, $\mathbf{J}$ has elements J_{ij} given by

$$J_{ij} = \frac{\partial f_i}{\partial x_j} \tag{31}$$

- For our current example we have

$$\mathbf{J} = \begin{bmatrix} \partial f_1/\partial x & \partial f_1/\partial y \\ \partial f_2/\partial x & \partial f_2/\partial y \end{bmatrix} = \begin{bmatrix} y\cos(xy) & x\cos(xy) \\ -6 & 2y \end{bmatrix}$$

- We can now derive the multi-dimensional Newton iteration, by considering a multivariate Taylor series expansion, paralleling what we did in the 1-d case:

$$\mathbf{0} = \mathbf{f}(\mathbf{x}^\star) = \mathbf{f}(\mathbf{x}^{(n)}) + \mathbf{J}[\mathbf{x}^{(n)}] \cdot (\mathbf{x}^\star - \mathbf{x}^{(n)}) + O((\mathbf{x}^\star - \mathbf{x}^{(n)})^2) \quad (32)$$

where the notation $\mathbf{J}[\mathbf{x}^{(n)}]$ means we evaluate the Jacobian matrix at $\mathbf{x} = \mathbf{x}^{(n)}$.

Dropping higher order terms, and replacing $\mathbf{x}^\star$ with $\mathbf{x}^{(n+1)}$, we have

$$\mathbf{0} = \mathbf{f}(\mathbf{x}^{(n)}) + \mathbf{J}[\mathbf{x}^{(n)}](\mathbf{x}^{(n+1)} - \mathbf{x}^{(n)}) \quad (33)$$

Defining $\delta\mathbf{x}^{(n)}$ via

$$\delta\mathbf{x}^{(n)} \equiv -(\mathbf{x}^{(n+1)} - \mathbf{x}^{(n)}) \tag{34}$$

the d-dimensional Newton iteration is given by

$$\mathbf{x}^{(n+1)} = \mathbf{x}^{(n)} - \delta\mathbf{x}^{(n)} \tag{35}$$

where the update vector, $\delta\mathbf{x}^{(n)}$, satisfies the $d \times d$ *linear system*

$$\mathbf{J}[\mathbf{x}^{(n)}]\,\delta\mathbf{x}^{(n)} = \mathbf{f}(\mathbf{x}^{(n)}) \tag{36}$$

- Again note that the Jacobian matrix, $\mathbf{J}[\mathbf{x}^{(n)}]$, has elements

$$J_{ij}[\mathbf{x}^{(n)}] = \left.\frac{\partial f_i}{\partial x_j}\right|_{\mathbf{x}=\mathbf{x}^{(n)}} \tag{37}$$

- At each step of the Newton iteration, the linear system (36) can, of course, be solved using an appropriate linear solver (e.g. general, tridiagonal, or banded).

General Structure of a Multidimensional Newton Solver

$\mathbf{x}$: Solution vector

$\mathbf{res}$: Residual vector

$\mathbf{J}$: Jacobian matrix

$\mathbf{dx}$: Update vector

$\mathbf{x} = \mathbf{x}^{(0)}$

do while $\|\mathbf{dx}\|_2 > \epsilon$

    do i = 1 , neq

        res(i) $= f_i(\mathbf{x})$

        do j = 1 , neq

            J(i,j) $= [\partial f_i / \partial x_j)](\mathbf{x})$

        end do

    end do

    $\mathbf{dx} =$ solve$(\mathbf{J}\ \mathbf{dx} = \mathbf{res})$

    $\mathbf{x} = \mathbf{x}$ - $\mathbf{dx}$

end do

Finite Difference Example: Non-Linear BVP

- Consider the nonlinear two-point boundary value problem

$$u(x)_{xx} + (uu_x)^2 + \sin(u) = F(x) \tag{38}$$

which is to be solved on the interval

$$0 \leq x \leq 1 \tag{39}$$

with the boundary conditions

$$u(0) = u(1) = 0 \tag{40}$$

- As we did for the case of the linear BVP, we will approximately solve this equation using $O(h^2)$ finite difference techniques. As usual we introduce a uniform finite difference mesh:

$$x_j \equiv (j-1)h \qquad j = 1, 2, \cdots N \qquad\qquad h \equiv (N-1)^{-1} \tag{41}$$

- Then, using the standard $O(h^2)$ approximations to the first and second derivatives

$$u_x(x_j) = \frac{u_{j+1} - u_{j-1}}{2h} + O(h^2) \quad (42)$$

$$u_{xx}(x_j) = \frac{u_{j+1} - 2u_j + u_{j-1}}{h^2} + O(h^2) \quad (43)$$

the discretized version of (38-40) is

$$\frac{u_{j+1} - 2u_j + u_{j-1}}{h^2} + (u_j)^2 \left[\frac{u_{j+1} - u_{j-1}}{2h}\right]^2 + \sin(u_j) - F_j = 0\,; \quad j = 2 \ldots N-1 \quad (44)$$

$$u_1 = 0 \quad (45)$$

$$u_N = 0 \quad (46)$$

Note that we have cast the discrete equations in the canonical form $\mathbf{f}(\mathbf{u}) = \mathbf{0}$

- In order to apply Newton's method to the algebraic equations (45-46), we must compute the Jacobian matrix elements of the system.

- We first observe that due to the "nearest-neighbor" couplings of the unknowns u_j via the approximations (42-43), the Jacobian matrix is *tridiagonal* in this case.

- For the *interior* grid points, $j = 2 \ldots N$, corresponding to rows $2 \ldots N$ of the matrix, we have the following non-zero Jacobian elements:

$$J_{j\,,j} = -\frac{2}{h^2} + 2u_j \left[\frac{u_{j+1} - u_{j-1}}{2h}\right]^2 + \cos(u_j) \tag{47}$$

$$J_{j\,,j-1} = \frac{1}{h^2} - (u_j)^2 \frac{u_{j+1} - u_{j-1}}{2h^2} \tag{48}$$

$$J_{j\,,j+1} = \frac{1}{h^2} + (u_j)^2 \frac{u_{j+1} - u_{j-1}}{2h^2} \tag{49}$$

- For the *boundary* points, $j = 1$ and $j = N$, corresponding to the first and last row, respectively, of $\mathbf{J}$, we have

$$J_{1,1} = 1 \tag{50}$$

$$J_{1,2} = 0 \tag{51}$$

and

$$J_{N,N} = 1 \tag{52}$$

$$J_{N,N-1} = 0 \tag{53}$$

- Note that these last expressions correspond to the "trivial" equations

$$f_1 = u_1 = 0 \tag{54}$$

$$f_N = u_N = 0 \tag{55}$$

which have associated residuals

$$r_1^{(n)} = u_1^{(n)} \tag{56}$$

$$r_N^{(n)} = u_N^{(n)} \tag{57}$$

- Observe that if we initialize $u_1^{(0)} = 0$ and $u_N^{(0)} = 0$, then we will *automatically* have $\delta u_1^{(n)} = \delta u_N^{(n)} = 0$, which will yield $u_1^{(n)} = 0$ and $u_N^{(n)} = 0$ as desired.

- This is an example of the general procedure we have seen previously for imposing Dirichlet conditions; namely the conditions are implemented as "trivial" (linear) equations (but it is, of course, absolutely crucial to implement them *properly* in this fashion!)

- *Testing procedure:* We adopt the same technique used for the linear BVP case—we *specify* $u(x)$, then compute the function $F(x)$ that is required to satisfy (38); $F(x)$ is then supplied as input to the code, and we ensure that as $h \to 0$ we observe second order convergence of the computed finite difference solution $\hat{u}(x)$ to the continuum solution $u(x)$.

- *Example:* Taking

$$u(x) = \sin(4\pi x) \equiv \sin(\omega x) \tag{58}$$

then

$$\begin{aligned} F(x) &= u_{xx} + (uu_x)^2 + \sin(u) \qquad (59) \\ &= -\omega^2 \sin(\omega x) + \omega^2 \sin^2(\omega x) \cos^2(\omega x) + \sin(\sin(\omega x)) \end{aligned}$$

- We note that due to the nonlinearity of the system, we will actually find *multiple* solutions, depending on how we initialize the Newton iteration; this is illustrated with the on-line code `nlbvp1d`.

3. Determining Good Initial Guesses: Continuation

- It is often the case that we will want to solve nonlinear equations of the form

$$\mathbf{N}(\mathbf{x}; \bar{\mathbf{p}}) = 0 \tag{60}$$

where we have adopted the notation $\mathbf{N}(\cdots)$ to emphasize that we *are* dealing with a nonlinear system. Here $\mathbf{x} = (x_1, x_2 \ldots x_d)$ is, as previously, a vector of unknowns, with $\mathbf{x} = \mathbf{x}^\star$ *a* solution of (60).

- The quantity $\bar{\mathbf{p}}$ in (60) is another vector, of length m, which enumerates any additional parameters (generally adjustable) that enter into the problem; these could include: coupling constants, rate constants, "perturbation" amplitudes etc.

- The nonlinearity of any particular system of the form (60) may make it *very* difficult to compute $\mathbf{x}^\star$ without a good initial estimate $\mathbf{x}^{(0)}$; in such cases, the technique of *continuation* often provides the means to generate such an estimate.

- *Continuation:* The basic idea underlying continuation is to "sneak up" on the solution by introducing an additional parameter, ϵ (the continuation parameter), so that by *continuously* varying ϵ from 0 to 1 (by convention), we vary *from*:

  1. A problem that we know how to solve, or for which we already have a solution.

  to

  2. The problem of interest.

• Schematically we can sketch the following picture:



• Note that we thus consider a *family* of problems

$$\mathbf{N}_\epsilon(\mathbf{x};\bar{\mathbf{p}}) = 0 \tag{61}$$

with corresponding solutions

$$\mathbf{x}_\epsilon = \mathbf{x}^\star_\epsilon \tag{62}$$

- The efficacy of continuation generally depends on two cruical points:

1. $\mathbf{N}_0(\mathbf{x}; \bar{\mathbf{p}})$ has a known or easily calculable root at $\mathbf{x}_0^\star$.

2. Can often choose $\Delta\epsilon$ judiciously (i.e. sufficiently small) so that

$$\mathbf{x}^\star_{\epsilon-\Delta\epsilon}$$

is a "good enough" initial estimate for

$$\mathbf{N}_\epsilon(\mathbf{x}; \bar{\mathbf{p}}) = 0$$

- Again, schematically, we have



where we note that we may have to adjust (adapt) $\Delta\epsilon$ as the continuation proceeds.

Continuation: Summary and Comments

- Solve sequence of problems with $\epsilon = 0, \epsilon_2, \epsilon_3 \ldots 1$ using previous solution as initial estimate for each $\epsilon \neq 0$.
- Will generally have to tailor idea on a case-by-case basis.
- Can often identify ϵ with one of the p_i (intrinsic problem parameters) *per se*.
- The first problem, $\mathbf{N}_0(\mathbf{x}, \bar{\mathbf{p}}) = 0$, can frequently be chosen to be *linear*, and therefore "easy" to solve, modulo sensitivity/poor conditioning.
- For time-dependent problems, *time evolution* often provides "natural" continuation; $\epsilon \rightarrow t$, and we can use $\mathbf{x}^\star(t - \Delta t)$ as the initial estimate $\mathbf{x}^{(0)}(t)$.